

Histone H3 (Phospho Ser10Thr11) Rabbit mAb (AR1860)

Key Features

Host Species:	Rabbit
Reactivity:	Human, Mouse, Rat
Applications:	WB, IHC, IF, IP, ELISA
Isotype:	IgG, Kappa
MW:	15kDa (Calculated) 17kDa (Observed)

Recommended Dilution Ratios

IHC:	1:200-1000
WB:	1:2000-10000
IF:	1:200-1000
ELISA:	1:5000-20000
IP:	1:50-200

Storage

-15°C to -25°C/1 year (Do not lower than -25°C)

Basic Information

Clonality	Monoclonal
-----------	------------

Immunogen Information

Specificity

Histone H3 (Phospho Ser10/Thr11) Monoclonal Antibody detects endogenous levels of Histone H3 protein only when phosphorylated at Ser10/Thr11. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):RKstGG

Target Information

Gene name HIST1H3A/HIST1H3B/HIST1H3C/HIST1H3D/HIST1H3E/HIST1H3F/HIST1H3G/HIST1H3H/HIST1H3I/HIST1H3J/HIST2H3A/HIST2H3C/HIST2H3D/H3F3A/H3F3B


Protein Name Histone H3.1/Histone H3.2/Histone H3.3

	Organism	Gene ID	UniProt ID
Human		8350; 8351; 8352; 8353; 8354; 8355; 8356; 8357; 8358; 8968	P68431; Q71DI3; P84243
	Mouse	319152; 15077; 15078	P68433
	Rat	291159; 100361558	Q6LEDO; P84245

Cellular Localization Nucleus. Chromosome

Tissue specificity Blood, Epithelium, Kidney, Lung, Ovary, Spleen, Uterus

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Histone H3 (Phospho Ser10/Thr11) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody.

Lane 1: Hela

Lane 2: Hela was treated with Nocodazole (50ng/ml) for 20 hours

Lane 3: NIH-3T3


Lane 4: NIH-3T3 was treated with Nocodazole (50ng/ml) for 20 hours

Lane 5: C6

Lane6: C6 was treated with Nocodazole (50ng/ml) for 20 hours

Predicted band size: 15kDa

Observed band size: 17kDa



Human testis was stained with anti-Histone H3 (Phospho Ser10/Thr11) rabbit antibody.

For Research Use Only